

# Talonavicular Joint Fusion

## INTRODUCTION

The talonavicular joint is located between the bone under the ankle (talus) and the navicular. It is responsible for some of the side-to-side movement of the hindfoot. At times it can become arthritic, resulting in pain, swelling and loss of function. Fusing the talonavicular joint involves joining the talus and navicular together, and is an effective way of decreasing pain and improving the function of the foot. After a talonavicular fusion most patients are able to walk without a limp and have considerably less pain.

## THE SURGERY

Talonavicular fusion surgery involves a number of steps:

- General anaesthetic, nerve block and IV antibiotics
- Tourniquet around the thigh
- Incision over the top of the foot
- Removal of any remaining cartilage
- Bone graft harvest (heel bone or distal tibia)
- Insertion of bone graft into the talonavicular joint to stimulate fusion
- Internal fixation with plates and screws
- Check x-rays
- Closure of incisions with sutures
- Plaster cast or surgical camboot (VACOCast)

## GUIDELINES FOR POST-OP RECOVERY

### HOSPITAL ADMISSION

- In hospital for 1 night, IV antibiotics

### FIRST 2 WEEKS

- Elevate foot and rest
- Strictly **NON-WEIGHTBEARING** in surgical boot
- Mobilise with knee scooter/crutches/frame
- Boot and dressings to stay dry and intact
- Strong painkillers as required
- Aspirin 100mg, Vitamin D 5000 units and Vitamin C 1g per day
- Antibiotics for 5 days
- Wiggle toes to encourage circulation

### 2 WEEK POST-OP APPOINTMENT

- Review by nurse & removal of dressings and sutures

### WEEKS 3-6

- Strictly **NON-WEIGHTBEARING** in boot
- Surgical boot on at ALL times, including in bed, except for rehab exercises & showers (seated with shower chair)
- Mobilise with knee scooter/crutches/frame
- Elevate foot when resting
- Continue Aspirin, Vitamin D and Vitamin C
- Podiatry/physiotherapy review (arrange first appointment 2-5 days after 2-week post-op check)
- **REHAB - PODIATRY/PHYSIOTHERAPY:**
  - Joint mobilisation, soft tissue manipulation, nerve desensitisation
  - Gentle active ankle plantarflexion/dorsiflexion and toe flexion/extension - **NO** inversion/eversion
  - Soft tissue massage and scar mobilisation
  - General upper body, core, hip and knee exercises

### 6 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

### WEEKS 7-12

- Surgical boot on at all times during the day except for rehab exercises & showers (with shower chair)
- Can sleep out of boot at night
- Elevate foot when resting
- Continue Aspirin, Vitamin D and Vitamin C
- **PODIATRY/PHYSIOTHERAPY:**
  - Commence partial weightbearing in surgical boot with crutches/frame, and progress up to full weightbearing as tolerated
  - Passive & active ROM ankle, midfoot, forefoot joints

### 12 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

### FROM 12 WEEKS

- **PODIATRY/PHYSIOTHERAPY:**
  - Transition out of boot into normal supportive shoes
  - Gait re-training, continue strength and conditioning

### 6 MONTH POST-OP APPOINTMENT

- CT scan to confirm fusion and review by Dr Zilko

Full recovery is usually 6-12 months.

*Every patient's recovery is individual and depends on the severity of the injury/disease and complexity of the surgery.*

Copyright 2024 Dr Simon Zilko. All rights reserved.